

This material has been produced and funded by Novo Nordisk for adult patients in the United Kingdom (UK) who have already been prescribed Wegovy® ▼ (semaglutide injection) for cardiovascular risk reduction only. This material is not a substitute for clinical advice provided by your healthcare professional nor is it a substitute for the Patient Information Leaflet, which you are advised to read in full. Please contact your healthcare professional if you have any questions about your treatment and for clinical advice. For further information about Wegovy®, please refer to the Patient Information Leaflet [HERE](#).

Getting started with Wegovy® ▼ (semaglutide injection)

Information about Wegovy® and cardiovascular risk reduction
in adults with a history of serious heart or circulatory disease*

*Serious heart or circulatory disease refers to heart attack, stroke, or poor blood flow to the legs.

▼ This medicine is subject to additional monitoring. This will allow quick identification of new safety information. **Reporting of side effects:** if you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in the package leaflet. You can also report side effects directly via the Yellow Card Scheme at <https://yellowcard.mhra.gov.uk>. Adverse events should also be reported to Novo Nordisk Limited (Telephone Novo Nordisk Customer Care Centre 0800 023 2573). Calls may be monitored for training purposes.

By reporting side effects, you can help provide more information on the safety of this medicine.

Please refer to the [Patient Information Leaflet](#) found in the product carton for

- further information on Wegovy®
- further information on how to use Wegovy®
- a full list of side effects, warnings and precautions.

Information on possible side effects for Wegovy® can be found on pages 15-16 of this material.

Information on Warnings and precautions for Wegovy® can be found on pages 12-14 of this material.

This material is designed to be viewed digitally.

It contains hyperlinks which are viewable online only. UK26SEMO00102 | Date of preparation: May 2026



This image does not represent a real patient





Welcome to Wegovy®

This is where you will find information, advice, and tips to getting started with Wegovy®. For further information, please visit wegovy.co.uk

Contents

Understanding cardiovascular disease	3
Understanding the need to reduce your risk of cardiovascular events when you already have serious heart or circulatory disease	4
How does Wegovy® work?	5
Get to know your Wegovy® FlexTouch® pen	6
How to use once-weekly Wegovy®	7
A pre-filled pen that you can administer yourself	8-9
Stepping up your dose	10
Missed doses, storage and disposal	11
Wegovy® safety information	12-14
Very common and common side effects	15
Serious side effects	16
Eating a balanced diet	17
Tips for being more active	18
PenCycle®	19



Understanding cardiovascular disease


Cardiovascular disease (CVD) is a term for a number of conditions that affect the heart or blood vessels and is typically associated with fatty deposit build up in the arteries including those in organs such as the heart, brain, kidneys and eyes.

In the UK, cardiovascular disease is one of the most common causes of death and disability.

However, making healthy lifestyle choices can help to prevent it.

Several factors are known to increase the risk of heart attacks and strokes, for example:

- High blood pressure
- Smoking
- High cholesterol
- Diabetes
- Living with overweight or obesity



Other risk factors that affect your risk of developing cardiovascular disease include:

- Being over 50 years old
- Being male (men are more likely to develop cardiovascular disease at an earlier age than women)
- Family history of cardiovascular disease
- Ethnicity (UK population of south Asian, Black African and African Caribbean backgrounds are more likely to have other risk factors for CVD)
- Having existing serious cardiovascular disease, for example a history of stroke, heart attack or poor circulation to the limbs, is itself a risk factor for further cardiovascular events – many patients experience recurrent cardiovascular events

Understanding the need to reduce your risk of cardiovascular events when you already have serious heart or circulatory disease



A healthy heart is essential for an active life, yet many of us face risks that can lead to serious conditions such as a heart attack or stroke. Understanding and managing these risks is crucial for maintaining your health. If you have already experienced a heart attack, stroke or poor blood flow to the limbs, you are at risk of having a serious cardiovascular event (heart-related death, another stroke or heart attack). It is important, therefore, that you manage these risks to help reduce the risk of another serious event.

You and your healthcare professional have decided to make Wegovy® part of your treatment plan in addition to diet and physical activity.

This brochure will help explain why, if you have a history of a heart attack, stroke, or poor blood flow to the limbs, reducing the risk of a further serious cardiovascular event is important. It will also give you information about a balanced diet and tips for being more active.



How does Wegovy® work?

Wegovy® is a medicine that contains the active substance semaglutide.

Your body

Your body sends signals (via appetite hormones) to the brain to tell you that you are hungry or crave certain foods.

Wegovy® for risk reduction of serious heart and circulatory issues

Wegovy® is used in addition to diet and physical activity to reduce the risk of serious heart issues (heart-related death, heart attacks, strokes) in adults with a history of heart disease (like a heart attack, stroke or poor blood flow to the legs) and either obesity or overweight (BMI ≥ 27 kg/m²).



Wegovy®

Similar to a natural hormone called glucagon-like peptide-1 (GLP-1) that is released from the intestine after a meal, Wegovy® works by acting on receptors in the brain that control your appetite, causing you to feel fuller and less hungry and experience less cravings for food. The exact mechanism by which Wegovy® reduces cardiovascular risk is not known.

This image does not represent a real patient

Get to know your Wegovy® FlexTouch® pen

Each Wegovy® pack consists of 1 pre-filled Wegovy® FlexTouch® pen which contains 4 of your prescribed doses of Wegovy®, to be used once a week for 4 weeks, 4 disposable needles and the Patient Information Leaflet. After having injected the 4 doses, there might still be solution left in the pen despite having administered correctly. Any solution left is insufficient for a dose and the pen should be disposed of.




This image is of the 2.4 mg FlexTouch® pen and is a representation only. It does not reflect the actual size of the pen.

How to use once-weekly Wegovy®

Always use this medicine exactly as your doctor has told you.
Check with your doctor, pharmacist or nurse if you are not sure.

Where to Inject

Wegovy® is given as an injection under the skin (subcutaneous injection). Do not inject it into a vein or muscle. Before you use the pen for the first time, ask your doctor or nurse on where to inject. The best places to give the injection are:



You may inject in the same body area each week, but make sure it is not in the same spot as used the last time.

When to Inject

Inject once a week and if possible, on the same day each week. This can be any time of the day – regardless of meals. If you need to, you can change the day of your weekly injection as long as it has been at least 3 days since your last injection. After selecting a new dosing day, continue with once a week dosing.



Try to get into a routine for using Wegovy® once a week.

For example, pick a specific day and time, like on a Sunday before you go to bed.

If you use more Wegovy® than you should, talk to your doctor straight away. You may get side effects such as feeling sick (nausea).

Do not stop using Wegovy® without talking to your doctor.


If you have any further questions on the use of this medicine, ask your doctor, pharmacist or nurse.

For full information on how to use your Wegovy® FlexTouch® pen, refer to the Patient Information Leaflet [HERE](#).

A pre-filled pen that you can administer yourself.

Before you use the pen for the first time, ask your doctor or nurse how to use it.

1. Get started



Check the name and dose of the pen to make sure it contains the prescribed dose of Wegovy®. Pull off the pen cap. **Check that the Wegovy® in the pen is clear and colourless. If Wegovy® looks cloudy or coloured, do not use the pen.**

2. Attach a needle



Always use a new needle for each injection, never use a bent or damaged needle. Check the paper tab and outer needle cap for damages. If you see any damage, dispose of it and use a new needle. Tear off the paper tab. Push the needle straight onto the pen. Turn until it is on tight. The needle is covered by two caps. Pull off the outer needle cap and keep it for later. Pull off the inner needle cap and dispose of it.

3. Check the Wegovy® flow



Check the flow before the first injection with each new pen. Turn the dose selector until the flow check symbol (■ ■ ▲) lines up with the dose pointer. Hold the pen with the needle pointing up. Press and hold in the dose button until the dose counter returns to -0-. A drop of Wegovy® should appear at the needle tip. This drop indicates that the pen is ready for use. If a drop does not appear, check the flow again. This should only be done twice. If there is still no drop, change the needle and check the flow once more. Do not use the pen if a drop of Wegovy® still does not appear.

A pre-filled pen that you can administer yourself.

Before you use the pen for the first time, ask your doctor or nurse how to use it.

4. Set the dose




Turn the dose selector until the dose counter stops, and it shows the prescribed dose. When the prescribed dose lines up with the dose pointer, the dose has been selected. Do not set the dose by counting the number of clicks you hear.

5. Inject the dose



Choose an injection site on your upper arms, upper legs or stomach (abdomen), keeping 5 cm away from your belly button. Insert the needle into your skin. Make sure the dose counter is visible. Press and hold down the dose button until the dose counter shows -0-. Keep pressing the dose button with the needle in your skin and slowly count to 6. The -0- must line up with the dose pointer, you may hear or feel a click when the dose counter returns to 0. Remove the needle from your skin. If blood appears at the injection site, press lightly on the area to stop the bleeding.

6. After injection



Lead the needle tip into the outer needle cap on a flat surface without touching the needle. Never try to put the inner cap back on the needle as you may stick yourself with the needle. Once the needle is covered, carefully push the outer needle cap completely on. Unscrew the needle. Always dispose of the needle immediately after each injection using a sharps bins. Never store your pen with the needle attached. Keep your sharps bins in a safe place so it's not a risk to other people and is out of the sight and reach of children. Sharps bins must not be disposed of in household waste. Discuss with your healthcare professional the procedure for disposing of sharps bins in your area. Put the pen cap on the pen after each use to protect Wegovy® from light.

Stepping up your dose


Always use this medicine exactly as your doctor has told you. Check with your doctor, pharmacist or nurse if you are not sure.

Your treatment will start at a low dose which will be gradually increased over 16 weeks to help reduce the likelihood of gastrointestinal-related side effects, such as feeling sick (nausea), being sick (vomiting), diarrhoea, constipation, and stomach pain.

When you first start using Wegovy®, the starting dose is 0.25 mg once-weekly. Your healthcare professional will instruct you to gradually increase your dose every 4 weeks until you reach the recommended maintenance dose of 2.4 mg once-weekly, or your maximum tolerated dose.

Once you reach the dose of 2.4 mg once-weekly, do not increase the dose further. 2.4 mg is the maximum maintenance dose licensed for cardiovascular risk reduction.

In case of significant gastrointestinal symptoms, your healthcare professional may consider a delay to increasing your dose or lower to the previous dose until your symptoms are better. It's important to follow the dosing schedule as recommended by your healthcare professional and discuss any side effects you may experience with them. Check with your healthcare professional if you are not sure.



Missed doses, storage and disposal

What to do if you miss a dose

- Do not take a double dose to make up for a missed one
- If you do miss a dose, try to remember the 5-day rule:
 - **If it is 5 days or less since you should have used Wegovy®:** Use it as soon as you remember, then inject your next dose as usual on your scheduled day
 - **More than 5 days since you should have used Wegovy®:** Skip the missed dose and inject as usual on your next scheduled day



How to store Wegovy®

- Keep your Wegovy® pen out of the sight and reach of children
- Do not freeze Wegovy® and do not use if it has been frozen
- Keep the pen cap on when the pen is not in use in order to protect it from light
- Do not use this medicine after the expiry date

Before opening: Keep the pen refrigerated (2°C to 8°C). Keep the pen away from the cooling element.

During use: You can keep the pen for 6 weeks when stored at a temperature below 30°C or in a refrigerator (2°C to 8°C) away from the cooling element.

Never store your pen with the needle attached.



How to dispose of your needles and pens

When the pen is empty, dispose of the pen without a needle on as instructed by your doctor, pharmacist, nurse or local authorities. You can now also recycle your pen through PenCycle®, see page 19 of this material for further information. The pen cap and the empty carton can be disposed of in your household waste.

Always dispose of the needle immediately after each injection using a sharps bin. Never store your pen with the needle attached. Keep your sharps bin in a safe place so it's not a risk to other people and is out of the sight and reach of children. Sharps bins must not be disposed of in household waste. Discuss with your healthcare professional the procedure for disposing of sharps bins in your area.



Wegovy® safety information

Please refer to the Patient Information Leaflet [HERE](#) for a full list of side effects, warnings and precautions.

Do not use Wegovy® if you are allergic to semaglutide or any of the other ingredients of this medicine: disodium phosphate dihydrate, propylene glycol, phenol, sodium hydroxide/hydrochloric acid (for pH adjustment), water for injection.

Warnings and precautions Talk to your doctor, pharmacist or nurse before using Wegovy® or during treatment if you have:

Effects on the digestive system

During treatment with Wegovy®, you may feel sick (nausea) or be sick (vomiting), or have diarrhoea. These side effects can cause dehydration (loss of fluids). It is important that you drink enough fluids to prevent dehydration. This is especially important if you have kidney problems. Talk to your doctor if you have any questions or concerns.

Inflammation of the pancreas

If you have ever had pancreatitis (inflammation of the pancreas) which may cause severe pain in the stomach and back which does not go away; see page 16 of this material for further information.

Diabetic eye disease (retinopathy)

Fast improvements in blood sugar control may lead to a temporary worsening of diabetic eye disease. If you have diabetic eye disease and experience eye problems while taking Wegovy®, talk to your doctor.

Sudden changes to your eyesight

If you notice a sudden loss of vision or rapidly worsening eyesight during treatment with semaglutide, urgently contact your doctor.

This may be caused by a very rare side effect called non-arteritic anterior ischaemic optic neuropathy (NAION). Your doctor will refer you for an eye examination by an ophthalmologist and you may have to stop treatment with Wegovy®.

Patients with delayed stomach emptying (gastroparesis)

If you have slow (delayed) stomach emptying (called gastroparesis), use of Wegovy® may lead to serious or severe gastrointestinal adverse events. Talk to your doctor before using Wegovy®.

▼ This medicine is subject to additional monitoring. This will allow quick identification of new safety information. **Reporting of side effects:** if you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in the package leaflet. You can also report side effects directly via the Yellow Card Scheme at <https://yellowcard.mhra.gov.uk>. Adverse events should also be reported to Novo Nordisk Limited (Telephone Novo Nordisk Customer Care Centre 0800 023 2573). Calls may be monitored for training purposes.

By reporting side effects, you can help provide more information on the safety of this medicine.

Wegovy® safety information

Please refer to the Patient Information Leaflet [HERE](#) for a full list of side effects, warnings and precautions.

Low blood sugar (hypoglycaemia)

Wegovy® can cause low blood sugar. The warning signs of low blood sugar may come on suddenly. They can include cold sweat, cool pale skin, headache, fast heartbeat, feeling sick (nausea) or very hungry, changes in vision, feeling sleepy or weak, feeling nervous, anxious or confused, difficulty concentrating or shaking. If you have diabetes and are taking a sulfonylurea or an insulin with Wegovy® the risk of getting low blood sugar levels (hypoglycaemia) might increase. Your doctor may ask you to test your blood sugar levels. This will help your doctor decide if the dose of the sulfonylurea or insulin needs to be changed to reduce the risk of low blood sugar.

Surgery requiring anaesthesia

If you know that you are due to have surgery where you will be under anaesthesia (sleeping), please tell your doctor that you are taking Wegovy® prior to your surgical procedure.

Wegovy® can slow the emptying of your stomach, increasing the risk of stomach contents (e.g. food and drink) entering into your airways and lungs during surgery or procedures whilst you are under general anaesthesia or deep sedation. Your doctor will need to assess whether any modifications to the pre-procedure instruction and anaesthetic technique may be required.

Other medicines and Wegovy®

Tell your doctor, pharmacist or nurse if you are using, have recently used or might use any other medicines.

In particular, tell your doctor, pharmacist or nurse if you are using medicines containing the following:

- Warfarin or other similar medicines taken by mouth to reduce blood clotting (oral anti-coagulants). When you start treatment with e.g. warfarin or similar medicines, frequent blood testing to determine the ability of your blood to clot may be required.

Pregnancy and breast-feeding

This medicine should not be used during pregnancy, as it is not known if it may affect your unborn child. Therefore, it is recommended to use contraception while using this medicine. If you wish to become pregnant, you should stop using this medicine at least two months in advance. If you become or are pregnant, think you may be pregnant or are planning to have a baby when using this medicine, talk to your doctor straight away, as your treatment will need to be stopped.

You should not use this medicine if you are breast-feeding, as it is unknown if it passes into breast milk.

▼ This medicine is subject to additional monitoring. This will allow quick identification of new safety information. **Reporting of side effects:** if you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in the package leaflet. You can also report side effects directly via the Yellow Card Scheme at <https://yellowcard.mhra.gov.uk>. Adverse events should also be reported to Novo Nordisk Limited (Telephone Novo Nordisk Customer Care Centre 0800 023 2573). Calls may be monitored for training purposes.

By reporting side effects, you can help provide more information on the safety of this medicine.

Wegovy® safety information

Please refer to the Patient Information Leaflet [HERE](#) for a full list of side effects, warnings and precautions.

Driving and using machines

Wegovy® is unlikely to affect your ability to drive and use machines. Some patients may feel dizzy when taking Wegovy® mainly during the first 3 months of treatment (see section 4 of the Patient Information Leaflet). If you feel dizzy you should not drive or operate machines until you feel better. If you need any further information, talk to your doctor, pharmacist or nurse.

People with diabetes

Tell your doctor if you have diabetes. Your doctor may adjust the dose of your diabetes medicines to prevent you from getting low blood sugar.

- Wegovy® must not be used as a substitute for insulin.
- Do not mix Wegovy® up with other medicines that you inject (e.g. insulins).
- Do not use Wegovy® in combination with other medicines that contain GLP-1 receptor agonists.

For diabetics using this medicine in combination with a sulfonylurea or insulin, low blood sugar (hypoglycaemia) may occur which may reduce your ability to concentrate.

Do not drive or use machines if you get any signs of low blood sugar. Talk to your doctor for further information.

Sodium content

This medicine contains less than 1 mmol sodium (23 mg) per dose, i.e. essentially 'sodium-free'.

▼ This medicine is subject to additional monitoring. This will allow quick identification of new safety information. **Reporting of side effects:** if you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in the package leaflet. You can also report side effects directly via the Yellow Card Scheme at <https://yellowcard.mhra.gov.uk>. Adverse events should also be reported to Novo Nordisk Limited (Telephone Novo Nordisk Customer Care Centre 0800 023 2573). Calls may be monitored for training purposes.

By reporting side effects, you can help provide more information on the safety of this medicine.

Very common and common side effects

Please refer to the Patient Information Leaflet [HERE](#) for a full list of side effects, warnings and precautions.

The very common side effects when using Wegovy® (which may affect more than 1 in 10 people) are: headache, feeling sick (nausea), being sick (vomiting), diarrhoea, constipation, feeling weak or tired, and stomach pain. These usually go away over time.

Common side effects (which may affect up to 1 in 10 people) are: feeling dizzy, upset stomach or indigestion, burping, gas (flatulence), bloating of the stomach, inflamed stomach ('gastritis') – the signs include stomach ache, feeling sick (nausea) or being sick (vomiting), reflux or heartburn – also called 'gastro-oesophageal reflux disease', gallstones, hair loss, injection site reactions, change in the way food or drink tastes, change in skin sensation, low blood sugar (hypoglycaemia) in patients with diabetes. Complications of diabetic eye disease (diabetic retinopathy) is a common serious side effect.

Tips to manage nausea:



Try to

- Eat smaller meals
- Stop eating when full
- Take sips of a cool drink



Avoid

- Eating fatty or fried foods
- Eating spicy foods with strong smells
- Drinking alcohol

This is general advice on avoiding nausea. Always check with your healthcare professional for further advice.

Serious side effects

Please refer to the Patient Information Leaflet [HERE](#) for a full list of side effects, warnings and precautions.

Common:

may affect up to 1 in 10 people

- Complications of diabetic eye disease (retinopathy). If you have diabetes you should inform your doctor if you experience eye problems, such as changes in vision, during treatment with this medicine.

Uncommon:

may affect up to 1 in 100 people

- Inflamed pancreas (acute pancreatitis) which could cause severe pain in the stomach and back which does not go away. This is a serious, potentially life-threatening condition. You should see a doctor **immediately** if you experience such symptoms.

Stop using this medicine and seek urgent medical help if you experience: Severe, persistent pain in the stomach area (abdomen), with or without nausea and vomiting. This could be a sign of acute pancreatitis, which is serious and potentially life-threatening.

- Kidney or bladder stones. Signs may include back or lower abdomen pain, difficulty in urination or change in colour of your urine.

Rare:

may affect up to 1 in 1,000 people

- Severe allergic reactions (anaphylactic reactions, angioedema). **You should seek immediate medical help** and inform your doctor straight away if you get symptoms such as breathing problems, swelling of face, lips, tongue, and/or throat with difficulty swallowing, wheezing, fast heartbeat, pale and cold skin, feeling dizzy or weak.
- Hip fractures.

Very Rare:

may affect up to 1 in 10,000 people

- A medical condition of the eye called non-arteritic anterior ischaemic optic neuropathy (NAION), which may cause loss of vision without any pain. You should urgently contact your doctor if you notice sudden or gradually worsening eyesight (see section 2 of Patient Information Leaflet: "Sudden changes to your eyesight").

Not Known:

frequency cannot be estimated from the available data

- Bowel obstruction. A severe form of constipation with additional symptoms such as stomach ache, bloating, vomiting, etc.

▼ This medicine is subject to additional monitoring. This will allow quick identification of new safety information. **Reporting of side effects:** if you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in the package leaflet. You can also report side effects directly via the Yellow Card Scheme at <https://yellowcard.mhra.gov.uk>. Adverse events should also be reported to Novo Nordisk Limited (Telephone Novo Nordisk Customer Care Centre 0800 023 2573). Calls may be monitored for training purposes.

By reporting side effects, you can help provide more information on the safety of this medicine.

Eating a balanced diet

Treatment with Wegovy® should be in combination with a balanced, reduced-calorie diet to help support your overall health. A healthy balanced diet can give you the energy you need to lead an active life and help you avoid diet-related illness.

Consider the following tips to help you make healthier choices:



The plate method:

Aim to fill half your plate with vegetables or salad, fill one quarter with healthy protein (such as lean meat, fish, egg, tofu or beans) and one quarter with carbohydrates (such as potatoes, wholegrain rice, chapatti, yam, pasta or bread).



Make a healthy choice the easy choice:

Think about the meals that you are going to need. Make sure to have healthy snacks available (such as vegetable sticks, a piece of fruit, yoghurt or a small handful of nuts) in case you get hungry between meals.



Establish regular mealtimes:

Plan ahead for the meals that you are going to need. Try to have three balanced meals that are evenly spaced throughout the day.



Cut down, don't cut out:

Reduce the amount of food that you eat that is high in fat or added sugar. Trying to reduce the amount of these foods will be more sustainable than cutting them out all together.

Have more often

- Grilled, baked, poached or steamed foods
- A wide range of fruits and vegetables
- Wholemeal or wholegrain starchy carbohydrates (such as wholemeal breads, wholewheat pasta or brown rice)
- Oils and spreads which are lower in saturated fats such as olive, vegetable, sunflower or rapeseed. Use sparingly as all fats are high in calories

Have less often

- Fried or deep-fried foods
- Foods high in salt or sodium such as processed meals, cheese, crisps
- High energy snack foods such as sweets, chocolates, cakes or biscuits
- Fats that are high in saturated fats such as butter, ghee or lard
- Larger portions than needed

Tips for being more active

Being active can help improve heart and overall health and should be used alongside your treatment with Wegovy®. You should aim for 150 minutes of moderate intensity physical activity per week as this will help lower the risk of many physical and mental health conditions and improve your energy levels.

Here are some simple ideas and tips for getting started and exploring new activities.

Tips for getting started



Start off slow to avoid injury



Find a level that's right for you – everyone's starting point will be different



Try to be active for 30 minutes at least 5 days a week



Consider breaking exercise up into smaller sessions – for example you could try 10 minutes, 3 times each day



When the activity starts to get a little easier, consider adding a couple of extra minutes to your activity each time



Stop exercising and seek medical help if you do not feel well or are in pain during exercise

Not sure what to explore?

Here are some ideas for getting more active in your daily routine:



Try standing up while talking on the phone



Try walking or biking to the shops instead of driving



Try taking up a new active hobby, such as swimming



If you work from home, try to go for a walk to replace your usual commute time

If you have been prescribed Wegovy® using a pre-filled pen you can now recycle your pens with PenCycle®.

What is PenCycle®?

We have developed a **bespoke process** that lets us recycle our pre-filled pens after use.

Pre-filled Novo Nordisk pens contain medicine and are with pen users wherever they go. Because of this, they are made from high-quality, long-lasting materials that are too good to waste. By finding new ways to reuse them, we can help put an end to the unsustainable use of Earth's natural resources.

To find out more simply go to www.pen-cycle.co.uk

How can I PenCycle®?

PenCycling could not be simpler and is **completely free of charge**.



To get started, collect a **PenCycle® return box** from your nearest participating pharmacy, or order some online on our website (where you can also find a full list of participating pharmacies) and then follow the instructions by going to <https://www.pen-cycle.co.uk/diabetes-and-weight-management/how-do-i-pencycle.html>

Where can I PenCycle®?

You can drop off your full return box at any participating pharmacy or return them to Novo Nordisk by post as every return box has a FREEPOST Royal Mail address printed on the back. Pharmacies in certain parts of the UK are not able to accept return boxes due to current regulations.

For more information go to <https://www.pen-cycle.co.uk/diabetes-and-weight-management/where-do-i-pencycle.html>

You return it
We recycle it



Does not represent real patients



This material is produced and funded by Novo Nordisk Limited, with registered offices at
3 City Place, Beehive Ring Road, Gatwick, West Sussex, RH6 0PA.

Wegovy is a trademark owned by Novo Nordisk A/S. Novo Nordisk® is a trademark owned by Novo Nordisk A/S.

Wegovy® is a registered trademark of Novo Nordisk A/S. FlexTouch® is a registered trademark of Novo Nordisk A/S.

Novo Nordisk is a registered trademark of Novo Nordisk A/S. 2026© Novo Nordisk A/S, Novo Allé, DK-2880, Bagsvaerd, Denmark.

**This material is the intellectual property of Novo Nordisk and should not be placed on third-party websites.
Novo Nordisk has a responsibility for ensuring that all its materials reflect the most current information.**

UK26SEMO00102 | Date of preparation: May 2026

